Network Systems
Science & Advanced
Computing

Biocomplexity Institute & Initiative

University of Virginia

Estimation of COVID-19 Impact in Virginia

May 26th, 2021

(data current to May 22nd – May 25th)
Biocomplexity Institute Technical report: TR 2021-064



BIOCOMPLEXITY INSTITUTE

biocomplexity.virginia.edu

About Us

- Biocomplexity Institute at the University of Virginia
 - Using big data and simulations to understand massively interactive systems and solve societal problems
- Over 20 years of crafting and analyzing infectious disease models
 - Pandemic response for Influenza, Ebola, Zika, and others



Points of Contact

Bryan Lewis brylew@virginia.edu

Srini Venkatramanan srini@virginia.edu

Madhav Marathe marathe@virginia.edu

Chris Barrett@virginia.edu

Model Development, Outbreak Analytics, and Delivery Team

Przemyslaw Porebski, Joseph Outten, Brian Klahn, Alex Telionis, Srinivasan Venkatramanan, Bryan Lewis, Aniruddha Adiga, Hannah Baek, Chris Barrett, Jiangzhuo Chen, Patrick Corbett, Stephen Eubank, Ben Hurt, Dustin Machi, Achla Marathe, Madhav Marathe, Mark Orr,

Akhil Peddireddy, Asal Pilehvari, Erin Raymond, James Schlitt, Anil Vullikanti, Lijing Wang,

James Walke, Andrew Warren, Amanda Wilson, Dawen Xie



Overview

• Goal: Understand impact of COVID-19 mitigations in Virginia

Approach:

- Calibrate explanatory mechanistic model to observed cases
- Project based on scenarios for next 4 months
- Consider a range of possible mitigation effects in "what-if" scenarios

Outcomes:

- Ill, Confirmed, Hospitalized, ICU, Ventilated, Death
- Geographic spread over time, case counts, healthcare burdens

Key Takeaways

Projecting future cases precisely is impossible and unnecessary. Even without perfect projections, we can confidently draw conclusions:


- Case rates in Virginia continue to decline with pace accelerating in some districts
- VA mean weekly incidence down to 4/100K from 6/100K, US down (8 from 10 per 100K)
- Vaccination rates have slowed considerably, but population immunity now estimated over 65%
- Projections show declining rate overall across Commonwealth
- Recent updates:
 - Minor updates to measured acceptance levels and adjusted vaccination scenarios
 - Additional Fall resurgence study scenario testing resilience of population
 - Limited waning of natural immunity included in fit and projections, also with seroprevalence update


The situation continues to change. Models continue to be updated regularly.

Situation Assessment



Case Rates (per 100k) and Test Positivity





https://data.cms.gov/stories/s/q5r5-gjyu

County level test positivity from RT-PCR tests.

Green: <5.0%

(or with <20 tests in past 14 days)

Yellow: 5.0%-10.0%

(or with <500 tests and <2000 tests/100k and

>10% positivity over 14 days

Red: >10.0%

(and not "Green" or "Yellow")




District Trajectories

Goal: Define epochs of a Health District's COVID-19 incidence to characterize the current trajectory

Method: Find recent peak and use hockey stick fit to find inflection point afterwards, then use this period's slope to define the trajectory

Hockey stick fit




Trajectory	Description	Weekly Case Rate (per 100K) bounds	# Districts (prev week)
Declining	Sustained decreases following a recent peak	below -0.9	31 (31)
Plateau	Steady level with minimal trend up or down	above -0.9 and below 0.5	2 (4)
Slow Growth	Sustained growth not rapid enough to be considered a Surge	above 0.5 and below 2.5	2 (0)
In Surge	Currently experiencing sustained rapid and significant growth	2.5 or greater	0 (0)



District Trajectories – last 10 weeks

Status	# Districts (prev week)
Declining	30 (31)
Plateau	4 (4)
Slow Growth	1 (0)
In Surge	0 (0)

Curve shows smoothed case rate (per 100K) Trajectories of states in label & chart box Case Rate curve colored by Reproductive




Estimating Daily Reproductive Number


May 24th Estimates

Region	Date Confirmed R _e	Date Confirmed Diff Last Week
State-wide	0.769	0.072
Central	0.821	0.019
Eastern	0.815	0.121
Far SW	0.805	0.054
Near SW	0.823	0.112
Northern	0.748	0.181
Northwest	0.688	-0.189

Methodology

- Wallinga-Teunis method (EpiEstim¹) for cases by confirmation date
- Serial interval: updated to discrete distribution from observations (mean=4.3, Flaxman et al, Nature 2020)
- Using Confirmation date since due to increasingly unstable estimates from onset date due to backfill







^{1.} Anne Cori, Neil M. Ferguson, Christophe Fraser, Simon Cauchemez. A New Framework and Software to Estimate Time-Varying Reproduction Numbers During Epidemics. American Journal of Epidemiology, Volume 178, Issue 9, 1 November 2013, Pages 1505–1512, https://doi.org/10.1093/aje/kwt133


Changes in Case Detection

Timeframe (weeks)	Mean days	% difference from overall mean
July (26-30)	6.2	-2%
Aug (31-34)	4.9	-22%
Sept (35-38)	4.5	-28%
Oct (39-43)	4.5	-29%
Nov (44-47)	4.5	-28%
Dec (48-49)	4.3	-32%
Jan (00-04)	4.0	-37%
Feb (05-08)	3.5	-45%
Mar (09-13)	3.6	-43%
Apr (14-18)	3.2	-49%
Overall (13 - 18)	6.3	<u></u>




Test positivity vs. Onset to Diagnosis






Vaccination Administration Slows




Regional Vaccine courses initiated per day:

- Total counts of first dose of vaccines across regions
- Recent rise due to opening of vaccinations to 12-16 year olds




Shipments have slowed with decreased demand


Vaccinations Shift to Younger Populations



Vaccine Acceptance Data Sources



Household Pulse Survey COVID-19 Vaccination Tracker



COVIDcast / Facebook Survey

https://covidcast.cmu.edu

Census Household Pulse Surveys (HPS)

https://www.census.gov/library/visualizations/interactive/household-pulse-survey-covid-19-vaccination-tracker.html




Vaccination Acceptance – Comparison of Sources


Measured acceptance varies across sources:

- **COVIDcast / Facebook (FB):** Both corrected and actual measurement
- FluVax: Acceptance levels in VA for influenza vaccine during 2019-20 flu season
- Household Pulse (HPS): Census administered survey, but with some time delay till release (most recent for fortnight ending May 10th, 2021)
- HPS has highest overall, while FluVax is expectedly the lowest
- Corrected HPS and COVIDcast are very similar for VA at the state level, and approach FluVax


Virginia Vaccine Acceptance levels



State by State correlation COVIDcast and HPS



State level correlation between all four




Vaccination Acceptance by Region

Corrections to surveys:

- Facebook administered survey is timely and broad, but biased by who accesses Facebook and answers the survey
- Correction approach:
 - Calculate an over-reporting fraction based on reported vaccinations compared to VDH administration data
 - Cross-validate coarse corrections against HPS survey at the state level and corrected in same manner

Region	COVIDcast corrected	COVIDcast uncorrected	VDH proportion vaccinated	COVIDcast reported vaccinated
Central	67%	78%	55%	82%
Eastern	58%	80%	51%	83%
Far SW	54%	70%	43%	74%
Near SW	58%	74%	51%	74%
Northern	75%	85%	65%	88%
Northwest	64%	82%	53%	84%



Grey Bar: Survey measured and corrected acceptance

Green Bar: Proportion of eligible population


administered a vaccine


Dots: Proportion administered at least one dose for

each county



Vaccine Acceptance in Virginia - COVIDcast






Data Source: https://covidcast.cmu.edu


Acceptance remains high:


- Proportion of Virginians that have already or would definitely or probably accept vaccination if offered today
- Survey respondents are reporting high levels of vaccination of ~80% reflecting bias of the mechanism

Virginia - Vaccine Hesitancy

- **Top reasons for hesitancy**: side effects, distrust (increasing), unnecessary (increasing)
- More likely to take if recommended by: doctors and friends
- **Reasons unnecessary**: Not serious, not high risk, or other










Vaccine Acceptance by Region- COVIDcast

Levels of Acceptance and potential acceptance in flux:

- Nearly all the "Definitely Yes" have been vaccinated, yet there are 5-10% remaining across the regions
- Northwest and Southwest (to lesser degree) see growth in "probably not", seemingly from "definitely not"

Unvaccinated Only

All Respondents



Data Source: https://covidcast.cmu.edu





SARS-CoV2 Variants of Concern

Emerging new variants will alter the future trajectories of pandemic and have implications for future control

- Emerging variants can:
 - Increase transmissibility
 - Increase severity (more hospitalizations and/or deaths)
 - Limit immunity provided by prior infection and vaccinations
- Genomic surveillance remains very limited
 - Challenges ability to estimate impact in US to date and estimation of arrival and potential impact in future







Outbreak Info



SARS-CoV2 Variants of Concern

Lineage B.1.1.7

Prevalence: Levels have rapidly risen, as anticipated, and now are plateauing at national level and in many states, seemingly in VA as well

Transmissibility: Estimated increase of 50% compared to previous variants. B.1.1.7's mutations aids its infection efficiency, and thus boosts its overall levels of viremia; <u>study from Public Health England</u> shows contacts of B.1.1.7 cases are more likely (50%) to test positive than contacts of non-B.1.1.7 patients

Severity: Increased viremia also appears to increase the risk of hospitalization (60%) and mortality (60%). <u>Danish</u> study shows B.1.1.7 to have a 64% higher risk of hospitalization, while <u>Public Health Scotland</u> studies showed a range of 40% to 60%; <u>Study in Nature</u> based on UK data estimates B.1.1.7 cases have 60% higher mortality


Lineage B.1.351


Prevalence: Levels have remained low, as this variant's transmissibility can't compete with B.1.1.7, however, as more of the population becomes immune it may gain an advantage


Immune Escape: Many studies show that convalescent sera from previously infected individuals does not neutralize B.1.351 virus well which is <u>predictive</u> of <u>protection</u>, however, <u>vaccine induced immunity</u> shows <u>signs</u> of <u>effectiveness</u>

Lineage <u>B.1.429/427</u> and <u>B.1.526</u> and subvariants

 Combined account for around 20% of circulating virus, share may be shrinking as B.1.1.7 outcompetes








SARS-CoV2 Variants of Concern

Lineage P.1

- Prevalence: Nationally at 10.7%, lower but increasing in VA at 5.2%
- New study estimates 17-32% of all infections in Manaus in 2021 were reinfections, which helps explain data from Brazil demonstrating P.1's continued dominance in Rio despite presence of B.1.1.7


Lineage B.1.617.2 and related subvariants

- Continues to drive outbreak in India and neighbors, with immeasurable numbers of cases surpassing healthcare capacities in many regions
- Categorized as <u>VoC by Public Health England</u>, WHO, expect CDC to follow
- Strain shows <u>continued growth in UK, Europe</u> and in US
- <u>Several studies</u> estimate B.1.617.2 to have 100% faster growth than B.1.1.7, and UK study suggests a 13% advantage over B.1.1.7
- More studies show limited <u>immune escape</u> similar to B.1.351, however, still suggest protection remains for vaccinated
- PHE study shows limited efficacy of Astra-Zeneca with only one dose, efficacy returns following 2nd dose




20

Other State Comparisons



- Nearly all states are declining
- Growth out west has slowed, recent reporting artifacts in some states perturb the otherwise calm picture


Virginia and her neighbors



- VA and neighbors are all declining with steady pace
- Most neighbors are now below 10/100K level




Race and Ethnicity cases per 100K



Rates per 100K of each Racial-Ethnic population by Health District


- Each Health District's Racial-Ethnic population is plotted by their Hospitalization and Case Rate
- Points are sized based on their overall population size (overlapping labels removed)
- Change in rates over the last 2 weeks



26-May-21

Recent Incidence Compared to Summer 2020


Recent Incidence Compared to Weekly Summer Mean by County Mean: 0.92; Median: 0.6; IQR: 0.34-1.18



26-May-21 23

Shifting Age-Distributions

Hospitalizations in VA





26-May-21 24

Zip code level weekly Case Rate (per 100K)

Case Rates in the last week by zip code

- Concentrations in Southwest, which was preceded by cluster of increased HCW rates last week
- Still some universities in top 10
- Some counts are low and suppressed to protect anonymity, those are shown in white








Risk of Exposure by Group Size and HCW prevalence

Case Prevalence in the last week by zip code used to calculate risk of encountering someone infected in a gathering of randomly selected people (group size 25)


- **Group Size**: Assumes 2 undetected infections per confirmed case (ascertainment rate from recent seroprevalence survey), and shows minimum size of a group with a 50% chance an individual is infected by zip code (eg in a group of 23 in Green Bay, there is a 50% chance someone will be infected)
- HCW prevalence: Case rate among health care workers (HCW) in the last week using patient facing health care workers as the denominator



Current Hot-Spots

Case rates that are significantly different from neighboring areas or model projections


- **Spatial**: SaTScan based hot spots compare clusters of zip codes with weekly case prevalence higher than nearby zip codes to identify larger areas with statistically significant deviations
- **Temporal**: The weekly case rate (per 100K) projected last week compared to observed by county, which highlights temporal fluctuations that differ from the model's projections



Social Vulnerability and Recent Vaccination Rates

Comparison of social vulnerability and vaccination rate in last 2 weeks by county

• Social Vulnerability: Each county's Social Vulnerability Index (CDC) compared with the level of vaccination




26-May-21 28

Social Vulnerability and Total Vaccination Rates

Comparison of social vulnerability and total vaccination rate since the start of vaccination

• Social Vulnerability: Each county's Social Vulnerability Index (CDC) compared with the level of vaccination



26-May-21 29

Model Update – Adaptive Fitting



Adaptive Fitting Approach

Each county fit precisely, with recent trends used for future projection


 Allows history to be precisely captured, and used to guide bounds on projections


Model: An alternative use of the same meta-population model, PatchSim

- Allows for future "what-if" Scenarios to be layered on top of calibrated model
- Eliminates connectivity between patches, to allow calibration to capture the increasingly unsynchronized epidemic

External Seeding: Steady low-level importation

- Widespread pandemic eliminates sensitivity to initial conditions
- Uses steady 1 case per 10M population per day external seeding







Using Ensemble Model to Guide Projections

Ensemble methodology that combines the Adaptive with machine learning and statistical models such as:


- Autoregressive (AR, ARIMA)
- Neural networks (LSTM)
- Kalman filtering (EnKF)

Weekly forecasts done at county level.

Models chosen because of their track record in disease forecasting and to increase diversity and robustness.

Ensemble forecast provides additional 'surveillance' for making scenario-based projections.

Also submitted to CDC Forecast Hub.




Seroprevalence updates to model design

Several seroprevalence studies provide better picture of how many actual infections have occurred

 CDC Nationwide Commercial Laboratory Seroprevalence Survey estimated 14.5% [12% – 18%] seroprevalence as of March 4th – 17th up from 10.5% a month earlier

These findings are equivalent to an ascertainment ratio of ~2x in the future, with bounds of (1.3x to 3x)

- Thus for 2x there are 2 total infections in the population for every confirmed case recently
- This measure now fully tracks the estimated ascertainment over time
- Uncertainty design has been shifted to these bounds (previously higher ascertainments as was consistent earlier in the pandemic were being used)



https://covid.cdc.gov/covid-data-tracker/#national-lab



Calibration Approach

- Data:
 - County level case counts by date of onset (from VDH)
 - Confirmed cases for model fitting
- Calibration: fit model to observed data and ensemble's forecast
 - Tune transmissibility across ranges of:
 - Duration of incubation (5-9 days), infectiousness (3-7 days)
 - Undocumented case rate (1x to 7x) guided by seroprevalence studies
 - Detection delay: exposure to confirmation (4-12 days)
 - Approach captures uncertainty, but allows model to precisely track the full trajectory of the outbreak
- **Project:** future cases and outcomes generated using the collection of fit models run into the future
 - Mean trend from last 7 days of observed cases and first week of ensemble's forecast used
 - Outliers removed based on variances in the previous 3 weeks
 - 2 week interpolation to smooth transitions in rapidly changing trajectories



COVID-19 in Virginia:

Dashboard Updated: 5/18/2021

Cases, Hospitalizations and Deaths					
Total Cases* 670,834 (New Cases: 378)^		Total Hospitalizations** 29,290		Total Deaths 11,042	

[†] VDH adopted the updated CDC COVID-19 confirmed and probable surveillance case definitions on August 27, 2020. Found

Ou	utbreaks
Total Outbreaks*	Outbreak Associated Cases
3,519	75,664

At least two (2) has comminded cases are required to classify an outsteak.		
Testing (PCR Only)		
Testing Encounters PCR Only*	Current 7-Day Positivity Rate PCR Only**	
7,285,852	3.2%	

^{*} PCR" refers to "Reverse transcriptase polymerase chain reaction laboratory testing.

^{**} Lab reports may not have been received yet. Percent positivity is not calculated for days with incomplete data.

Multisystem Inflammatory Syndrome in Children		
Total Cases*	Total Deaths	
70	0	

^{*}Cases defined by CDC HAN case definition: https://emergency.cdc.gov/han/2020/han00432.as

Accessed 9:00am May 26, 2021

https://www.vdh.virginia.gov/coronavirus/

^{**} Hospitalization of a case is captured at the time VDH performs case investigation. This underrepresents the total number of hospitalizations in

New cases represent the number of confirmed and probable cases reported to VDH in the past 24 hours


Scenarios – Transmission Conditions


- Variety of factors continue to drive transmission rates
 - Seasonal impact of weather patterns, travel and gatherings, fatigue and premature relaxation of infection control practices
- Plausible levels of transmission can be bounded by past experience
 - Assess transmission levels at the county level from May 1, 2020 Sept 1, 2020 or current, whichever is highest
- Projection Scenario:
 - Adaptive: Control remains as is currently experienced into the future
 - Fatigued Control:
 - Highest level of transmission (95th percentile) increased by additional 5%
 - Transition to this level over 4 weeks, remain at this level for the summer, then return to Adaptive
- Additional study scenarios with Fall Resurgence:
 - Fall: Resurgence to worst of Fall 2020 starting in September and ramping up quickly



Scenarios – Mixed Variants Condition

- Variant B.1.1.7 has reached dominance in Virginia but no longer is growing at a predictable pace as variants compete
- Other variants with differing levels of transmissibility, immune escape, and impacts on severity also exist, to varying degrees
- Transmissibility boosting effects and growth of variants are no longer predictable
 - Rely on adaptive fitting to find transmissibility trends for projection
- Immune Escape
 - Many variants demonstrate the ability to evade immunity, both natural and vaccine-induced, uncertainty remains high thus this is not factored into the model
- Impact on Severity
 - Assume current variant prevalence remain relatively stable into future
 - Severity increases:
 - B.1.1.7= 1.6 times hospitalization and death
 - P.1 = 1.5 times hospitalization and death
 - B.1.351 = 1.5 times hospitalization and death










Scenarios – Vaccination Conditions

Assumed vaccine efficacies

- Pfizer/Moderna: 50% after first dose, 95% after second dose (3.5 week gap)
- J & J: 67% efficacy after first (and only) dose
- Delay to efficacy from doses is 14 days, immunity lasts at least 7m (NEJM study)


Two Vaccine Administration Scenarios

- Acceptance at county level = regional acceptance +/- relative current vax
- Status quo (no label): COVIDcast corrected acceptance estimate [VA ~65%] by Labor Day. County level heterogeneity based on cumulative uptake
- Optimistic (VaxOpt): Expand VA acceptance to 75% (~13% higher per county).
- Front-loaded rollout (two-thirds of the remaining in half the time)






	status quo	VaxOpt
Date		
2020-12-31	114.3K	114.3K
2021-01-31	768.0K	768.0K
2021-02-28	1.3M	1.3M
2021-03-31	2.6M	2.6M
2021-04-30	3.8M	3.8M
2021-05-31	4.2M	4.3M
2021-06-30	4.5M	4.9M
2021-07-31	4.8M	5.3M
2021-08-31	4.9M	5.6M
2021-09-30	4.9M	5.6M



Source: https://ckelly17.github.io/vaccine_dashboard.html

Weekly VA doses administered by manufacturer



Scenarios – Combined Conditions

Name	Txm Controls	Variant Boosting	Vax	Description
Adaptive	С	Mixed	SQ	Likely trajectory based on conditions remaining similar to how they are now
Adaptive-FatigueControl	F	Mixed	SQ	Worst case trajectory if control conditions deteriorate to highest transmission rates of the past
Adaptive-VaxOpt	С	Mixed	VO	Likely trajectory based on conditions remaining similar to now, but with improvements to vaccine acceptance
Adaptive-FatigueControl- VaxOpt	F	Mixed	VO	Worst case trajectory if control conditions deteriorate to worst of the past, with improvements to vax acceptance

Transmission Controls: C = Current levels persist into the future

F = Fatiguing controls drift to worst levels of last summer and persist

Variant Boosting: Mixed = Variety of variants, no future txm boosting, but with severity impacts from current levels

Vaccinations: SQ = Status quo acceptance leads to low rates of vaccination through the summer

VO = Vaccination acceptance reads to low rates of racematical through the se

VO = Vaccination acceptance optimistically expands with increased rates through the summer




Model Results




Outcome Projections


Confirmed cases



Estimated Hospital Occupancy




Daily Deaths



data, most recent dates are not complete


Daily Hospitalized






District Level Projections: Adaptive

Projections by Region




District Level Projections: Adaptive-VaxOpt

Projections by Region




District Level Projections: Adaptive-FatigueControl

Projections by Region



District Level Projections: Adaptive-FatigueControl-VaxOpt


Projections by Region




Hospital Demand and Bed Capacity by Region

Capacities* by Region – Adaptive-FatigueControl-DominantB117

COVID-19 capacity ranges from 80% (dots) to 120% (dash) of total beds





https://nssac.bii.virginia.edu/covid-19/vmrddash/

If Adaptive-FatigueControl-DominantB117 scenario persists:

No capacity challenges for hospital beds in near-term

UNIVERSITY OVIRGINIA

BIOCOMPLEXITY INSTITUTE


Resilience to Potential Fall Surge

Impact of expanded vaccine acceptance against a Fall Surge

- To further test Virginia's resilience to new transmission surges, implement a Fall Surge
 - Highest rate from Fall 2020, with a 2 week ramp up starting on Sept 1st 2021
- More highly vaccinated population more resilient to Fall Surge, with nearly 10k cases averted
- In worst case of sustained Fatigue
 Control 60K+ cases averted


Daily Infections

Virginia Daily Confirmed - Comparison



Cumulative Infections

Virginia Cumulative Confirmed - Comparison




26-May-21 46


Virginia's Progress on Population Immunity


Natural Immunity and Vaccines combine to produce a population level of immunity

- Duration of immunity from infection with SARS-CoV2 still not well understood
 - We assume a conservative 6 month period of protection for these calculations
 - Natural immunity is well calibrated to recent seroprevalence surveys
- Vaccine induced immunity is likely to last longer, we assume indefinite protection
 - This also assumes that all administered vaccines remain protective against current and future novel variants
- Population immunity depends on a very high proportion of the population getting vaccinated
 - Using regional vaccine acceptance




Region	% immune (est.)*
Central	68%
Eastern	63%
Far SW	63%
Near SW	67%
Northern	68%
Northwest	70%
Virginia	67%





Population immunity




COVID-19 Scenario Modeling Hub

Collaboration of multiple academic teams to provide national and state-by-state level projections for 4 aligned scenarios that vary vaccine rates (high – low) and levels of control (moderate and low)

Round 5 updates now available

Round 4 Results were published May 5th, 2021 in MMWR



26-May-21

Key Takeaways

Projecting future cases precisely is impossible and unnecessary. Even without perfect projections, we can confidently draw conclusions:

- Case rates in Virginia continue to decline with pace accelerating in some districts
- VA mean weekly incidence down to 4/100K from 6/100K, US down (8 from 10 per 100K)
- Vaccination rates have slowed considerably, but population immunity now estimated over 65%
- Projections show declining rate overall across Commonwealth
- Recent updates:
 - Minor updates to measured acceptance levels and adjusted vaccination scenarios
 - Additional Fall resurgence study scenario testing resilience of population
 - Limited waning of natural immunity included in fit and projections, also with seroprevalence update
- The situation continues to change. Models continue to be updated regularly.



References

Venkatramanan, S., et al. "Optimizing spatial allocation of seasonal influenza vaccine under temporal constraints." *PLoS Computational Biology* 15.9 (2019): e1007111.

Arindam Fadikar, Dave Higdon, Jiangzhuo Chen, Bryan Lewis, Srinivasan Venkatramanan, and Madhav Marathe. Calibrating a stochastic, agent-based model using quantile-based emulation. SIAM/ASA Journal on Uncertainty Quantification, 6(4):1685–1706, 2018.

Adiga, Aniruddha, Srinivasan Venkatramanan, Akhil Peddireddy, et al. "Evaluating the impact of international airline suspensions on COVID-19 direct importation risk." *medRxiv* (2020)

NSSAC. PatchSim: Code for simulating the metapopulation SEIR model. https://github.com/NSSAC/PatchSim

Virginia Department of Health. COVID-19 in Virginia. http://www.vdh.virginia.gov/coronavirus/

Biocomplexity Institute. COVID-19 Surveillance Dashboard. https://nssac.bii.virginia.edu/covid-19/dashboard/

Google. COVID-19 community mobility reports. https://www.google.com/covid19/mobility/

Biocomplexity page for data and other resources related to COVID-19: https://covid19.biocomplexity.virginia.edu/



Questions?

Points of Contact

Bryan Lewis brylew@virginia.edu

Srini Venkatramanan srini@virginia.edu

Madhav Marathe marathe@virginia.edu

Chris Barrett@virginia.edu

Biocomplexity COVID-19 Response Team

Aniruddha Adiga, Abhijin Adiga, Hannah Baek, Chris Barrett, Golda Barrow, Richard Beckman, Parantapa Bhattacharya, Jiangzhuo Chen, Clark Cucinell, Patrick Corbett, Allan Dickerman, Stephen Eubank, Stefan Hoops, Ben Hurt, Ron Kenyon, Brian Klahn, Bryan Lewis, Dustin Machi, Chunhong Mao, Achla Marathe, Madhav Marathe, Henning Mortveit, Mark Orr, Joseph Outten, Akhil Peddireddy, Przemyslaw Porebski, Erin Raymond, Jose Bayoan Santiago Calderon, James Schlitt, Samarth Swarup, Alex Telionis, Srinivasan Venkatramanan, Anil Vullikanti, James Walke, Andrew Warren, Amanda Wilson, Dawen Xie



Supplemental Slides




Estimating Daily Reproductive Number


May 8th Estimates

Region	Date of Onset R _e	Date Onset Diff Last Week
State-wide	0.587	-0.038
Central	0.643	-0.040
Eastern	0.602	-0.073
Far SW	0.714	0.053
Near SW	0.658	-0.052
Northern	0.581	-0.061
Northwest	0.701	0.039
Methodology		

Methodology

- Wallinga-Teunis method (EpiEstim¹) for cases by confirmation date
- Serial interval: updated to discrete distribution from observations (mean=4.3, Flaxman et al, Nature 2020)
- Using Confirmation date since due to increasingly unstable estimates from onset date due to backfill
- 1. Anne Cori, Neil M. Ferguson, Christophe Fraser, Simon Cauchemez. A New Framework and Software to Estimate Time-Varying Reproduction Numbers During Epidemics. American Journal of Epidemiology, Volume 178, Issue 9, 1 November 2013, Pages 1505–1512, https://doi.org/10.1093/aje/kwt133





Weekly Cases and Hospitalizations

Weekly confirmed cases

	Week Ending	Adaptive	Adaptive- Fatigued Control	Adaptive- VaxOpt	Adaptive- Fatigued Control -VaxOpt		-
	4/25/21	9,598	9,599	9,597			
	5/2/21	9,747	9,767	9,850			ľ
	5/9/21	,	8,894	9,683			<u> </u>
		8,048	8,086	10,150			
	5/23/21	7,392	7,428	11,250			
		6,713	6,561	13,152			
	6/6/21	5,994	5,553	16,388			
	6/13/21	5,365	4,521	21,163	UN:	VERSITY of VIR	C
.6	6/20/21	4,746	3,508	27,038	BIOCOMP	LEXITY IN	S


Weekly Hospitalizations

Week Ending	Adaptive- DominantB 117	Adaptive- BestPast- Dominant B117	Adaptive- Fatigued Control - DominantB1 17	
4/25/2				
1	750	750	750	
5/2/21	659	659	666	
5/9/21	548	548	601	
5/16/2	451	452	575	
5/23/2 1	373	372	580	
5/30/2 1	302	292	609	
6/6/21	241	219	654	
6/13/2	1,96	159	707	5/
6/20/2				

Agent-based Model (ABM)


EpiHiper: Distributed network-based stochastic disease transmission simulations

- Assess the impact on transmission under different conditions
- Assess the impacts of contact tracing



Synthetic Population

- Census derived age and household structure
- Time-Use survey driven activities at appropriate locations



Detailed Disease Course of COVID-19


- Literature based probabilities of outcomes with appropriate delays
- Varying levels of infectiousness
- Hypothetical treatments for future developments



ABM Social Distancing Rebound Study Design


Study of "Stay Home" policy adherence

- Calibration to current state in epidemic
- Implement "release" of different proportions of people from "staying at home"



Calibration to Current State

- Adjust transmission and adherence to current policies to current observations
- For Virginia, with same seeding approach as PatchSim



Impacts on Reproductive number with release

- After release, spike in transmission driven by additional interactions at work, retail, and other
- At 25% release (70-80% remain compliant)
- Translates to 15% increase in transmission, which represents a 1/6th return to pre-pandemic levels

